

## Year 1 – Week 13 Exam Questions

Question	1	2	3	4	5	6	7	8	Total
Marks									
Max Marks	5	4	5	6	4	6	7	8	45

*Suggested time: 55 minutes*

### Question 1

**In this question you must show detailed reasoning.**

(i) Express  $3^{\frac{7}{2}}$  in the form  $a\sqrt{b}$ , where  $a$  is an integer and  $b$  is a prime number. [2]

(ii) Express  $\frac{\sqrt{2}}{1-\sqrt{2}}$  in the form  $c + d\sqrt{e}$ , where  $c$  and  $d$  are integers and  $e$  is a prime number. [3]

### Question 2

(i) The equation  $x^2 + 3x + k = 0$  has repeated roots. Find the value of the constant  $k$ . [2]

(ii) Solve the inequality  $6 + x - x^2 > 0$ . [2]

### Question 3

The probability distribution of a random variable  $X$  is given in the table.

$x$	0	2	4	6
$P(X = x)$	$\frac{3}{8}$	$\frac{5}{16}$	$4p$	$p$

(i) Find the value of  $p$ . [2]

(ii) Two values of  $X$  are chosen at random. Find the probability that the product of these values is 0. [3]

### Question 4

In triangle  $ABC$ ,  $AB = 20$  cm and angle  $B = 45^\circ$ .

(i) Given that  $AC = 16$  cm, find the two possible values for angle  $C$ , correct to 1 decimal place. [4]

(ii) Given instead that the area of the triangle is  $75\sqrt{2}$  cm<sup>2</sup>, find  $BC$ . [2]

### Question 5

(i) The curve  $y = \frac{2}{3+x}$  is translated by four units in the positive  $x$ -direction. State the equation of the curve after it has been translated. [2]


(ii) Describe fully the single transformation that transforms the curve  $y = \frac{2}{3+x}$  to  $y = \frac{5}{3+x}$ . [2]

### Question 6

- (i) Express  $4x^2 - 12x + 11$  in the form  $a(x+b)^2 + c$ . [3]
- (ii) State the number of real roots of the equation  $4x^2 - 12x + 11 = 0$ . [1]
- (iii) Explain fully how the value of  $r$  is related to the number of real roots of the equation  $p(x+q)^2 + r = 0$  where  $p, q$  and  $r$  are real constants and  $p > 0$ . [2]

### Question 7

$OABC$  is a parallelogram with  $\overrightarrow{OA} = \mathbf{a}$  and  $\overrightarrow{OC} = \mathbf{c}$ .  $P$  is the midpoint of  $AC$ .



- (i) Find the following in terms of  $\mathbf{a}$  and  $\mathbf{c}$ , simplifying your answers.
- (a)  $\overrightarrow{AC}$  [1]
- (b)  $\overrightarrow{OP}$  [2]
- (ii) Hence prove that the diagonals of a parallelogram bisect one another. [4]

### Question 8



A particle  $P$  is moving along a straight line with constant acceleration. Initially the particle is at  $O$ . After 9 s,  $P$  is at a point  $A$ , where  $OA = 18$  m (see diagram) and the velocity of  $P$  at  $A$  is  $8 \text{ m s}^{-1}$  in the direction  $\overrightarrow{OA}$ .

- (i) (a) Show that the initial speed of  $P$  is  $4 \text{ m s}^{-1}$ . [2]
- (b) Find the acceleration of  $P$ . [2]
- $B$  is a point on the line such that  $OB = 10$  m, as shown in the diagram.
- (ii) Show that  $P$  is never at point  $B$ . [4]