

Year 1 – Week 10 Exam Questions – Mark Schee

Q1 Solution link <https://www.youtube.com/watch?v=0lqSSD9jYts>

Q2 Solution link <https://www.youtube.com/watch?v=Fn79BWQn-UE>

<https://www.youtube.com/watch?v=BYcuPM6bEIQ>

Q3Solution link <https://www.youtube.com/watch?v=rYYOO937nvY>

Mark Scheme

Question 1

(a)	$30^2 = 2a.300$ $a = 1.5$	M1 A1 (2)
(b)	$0^2 = 30^2 - 2 \times 1.25s$ $s = 360$ $300 + 30T + 360 = 1500$ $T = 28$	OR $0 = 30 - 1.25t_2$ $t_2 = 24$ $\frac{(20 + T + 24 + T)}{2} \times 30 = 1500$ $T = 28$ M1 A1 M1 A1 A1 (5)

Question 2

(a)	$v^2 = u^2 + 2as \Rightarrow 28^2 = u^2 + 2 \times 9.8 \times 17.5$ Leading to $u = 21$ *	cso	M1 A1 A1 (3)
(b)	$s = ut + \frac{1}{2}at^2 \Rightarrow 19 = 21t - 4.9t^2$ $4.9t^2 - 21t + 19 = 0$ $t = \frac{21 \pm \sqrt{21^2 - 4 \times 4.9 \times 19}}{9.8}$ $t = 2.99$ or 3.0 $t = 1.30$ or 1.3		M1 A1 DM1 A1 A1 (5)

Question 3

(a)	$240 = \frac{1}{2}(u + 34)10$	M1 A1
	$u = 14$	A1
		(3)
(b)	$34 = 14 + 10a \Rightarrow a = 2$	M1 A1
	$120 = 14t + \frac{1}{2} \times 2 \times t^2$	M1 A1
	$t^2 + 14t - 120 = 0$	
	Solving, $t = -20$ or 6	DM1
	$t = 6$	A1
	OR	
	$34 = 14 + 10a \Rightarrow a = 2$	M1 A1
	$v^2 = 14^2 + 2 \times 2 \times 120 \Rightarrow v = 26$	
	AND $26 = 14 + 2t$	M1 A1
	$t = 6$	DM1 A1
		(6)
		[9]

Question 4

Attempt to set up an equation of motion for 'A to B' using $s = 22$ and $t = 2$.	M1	<p>From A to B:</p> $s = 22 \text{ m}$ $u = u \quad 22 = 2u + \frac{1}{2}a(2)^2$ $v = ? \quad 22 = 2u + 2a \quad (1)$ $a = a$ $t = 2a$ <p>From A to C:</p> $s = 126 \text{ m}$ $u = u \quad 126 = 6u + \frac{1}{2}a(6)^2$ $v = ? \quad 126 = 6u + 18a$ $a = a \quad 21 = u + 3a \quad (2)$ $t = 6a$ <p>Solving (1) and (2) simultaneously:</p> $u = 6 \text{ ms}^{-1}, \quad a = 5 \text{ ms}^{-2}$
Sets up an equation in a and u only. (or a and V , where V is the velocity at B).	M1	
Obtains $22 = 2u + 2a$ or $22 = 2V - 2a$ (OE)	A1	
Sets up a second equation of motion using either 'A to C' or 'B to C' using either $s = 126$ and $t = 6$ (A to C) or $s = 104$ and $t = 4$ (B to C)	M1	
Obtains $126 = 6u + 18a$ or $104 = 4V + 8a$ (OE)	A1	
Attempts to solve the simultaneous equations. Note: If V is found, an attempt to find u must also be made.	M1	
Obtains correct values for u and a .	A1	
	7 marks	

Question 5

(a)	Correct use of distance formula with at least three terms correct. <i>Note: Omission of square root scores M0A0</i>	M1	$AB = \sqrt{(-2-6)^2 + (7-1)^2}$ $AB = 10$
	Correct Answer	A1	
		2 marks	
(b)	Correct use of $\frac{y_2 - y_1}{x_2 - x_1}$ (condone one sign error)	M1	$m = \frac{7-1}{-2-6} = -\frac{3}{4}$
	Correct answer (OE)	A1	
		2 marks	
(c)	Attempt to find the gradient by rearranging $4x - 3y - 10 = 0$ to make y the subject.	M1	$4x - 3y - 10 = 0$ $3y = 4x - 10$ $y = \frac{4}{3}x - \frac{10}{3}$ $\therefore \text{Gradient is } \frac{4}{3}$ $-\frac{3}{4} \times \frac{4}{3} = -1$ $\therefore \text{The lines are perpendicular.}$
	Attempt to use $m \times m_{\perp} = -1$ for <i>their</i> gradients. (may be implied by attempt to use of negative reciprocal)	B1ft	
	Reaches correct conclusion (the lines are perpendicular) from correct working.	A1	
		3 marks	

Question 6

Equates the two expressions.	M1	$y = 2x^2 - kx \quad y = x^2 - k$ $\therefore 2x^2 - kx = x^2 - k$ $x^2 - kx + k = 0$ The curves intersect provided that $\Delta \geq 0$ $\therefore (-k)^2 - 4(1)(k) \geq 0$ $k^2 - 4k \geq 0$ $k(k-4) \geq 0$ $\therefore \text{Solutions exist provided that } k \leq 0 \text{ or } k \geq 4$
Obtains $x^2 - kx + k = 0$	A1	
Attempt to calculate discriminant for <i>their</i> three-term quadratic.	M1	
Determines that the curves intersect provided that $\Delta \geq 0$ and obtains $k^2 - 4k \geq 0$	R1	
Attempt to use valid method for solving <i>their</i> quadratic inequality e.g. sketching the graph / table of values	M1	
Correct final answer from correct reasoning.	A1	
	6 marks	

