

Year 1 – Week 13 Exam Questions

Mark Scheme

Question 1

$m = -3$ (1 mark)

Use of $y - y_1 = m(x - x_1)$ with $(3, 1)$ or $(4, -2)$ (1mark)

$y = -3x + 10$ o.e. (1mark)

N.B. Answer left in the form $(y - 1) = -3(x - 3)$ or $(y - (-2)) = -3(x - 4)$ is awarded M1A1A0 as answers should be simplified by constants being collected

*i.e. scores
2 out of 3*

Question 2

Attempts $\overline{AB} = \overline{OB} - \overline{OA}$ or similar	M1
$\overline{AB} = 5i + 10j$	A1
	(2)
Finds length using 'Pythagoras' $ AB = \sqrt{(5)^2 + (10)^2}$	M1
$ AB = 5\sqrt{5}$	A1ft
	(2)


Question 3

(a)	States or uses $f(+3) = 0$	M1
	$4(3)^3 - 12(3)^2 + 2(3) - 6 = 108 - 108 + 6 - 6 = 0$ and so $(x - 3)$ is a factor	A1
		(2)
(b)	Begins division or factorisation so x $4x^3 - 12x^2 + 2x - 6 = (x - 3)(4x^2 + \dots)$	M1
	$4x^3 - 12x^2 + 2x - 6 = (x - 3)(4x^2 + 2)$	A1
	Considers the roots of their quadratic function using completion of square or discriminant	M1
	$(4x^2 + 2) = 0$ has no real roots with a reason (e.g. negative number does not have a real square root, or $4x^2 + 2 > 0$ for all x So $x = 3$ is the only real root of $f(x) = 0$ *	A1*
		(4)

Question 4

Realises that $k = 0$ will give no real roots as equation becomes $3 = 0$ (proof by contradiction)	B1
(For $k \neq 0$) quadratic has no real roots provided $b^2 < 4ac$ so $16k^2 < 12k$	M1
$4k(4k - 3) < 0$ with attempt at solution	M1
So $0 < k < \frac{3}{4}$, which together with $k = 0$ gives $0 \leq k < \frac{3}{4}$ *	A1*

Question 5

13(a)	$x^3 + 10x^2 + 25x = x(x^2 + 10x + 25)$	M1	
	$= x(x+5)^2$	A1	
		(2)	
(b)		M1	
		<p>A cubic with correct orientation</p> <p>Curve passes through the origin $(0, 0)$ and touches at $(-5, 0)$ (see note below for ft)</p>	A1ft
		(2)	
(c)	Curve has been translated a to the left	M1	
	$a = -2$	A1ft	
	$a = 3$	A1ft	
		(3)	

Question 6

Using distance = total area under graph (e.g. area of rectangle + triangle or trapezium or rectangle – triangle)	M1
e.g. $D = UT + \frac{1}{2} Th$, where h is height of triangle	A1
Using gradient = acceleration to substitute $h = aT$	M1
$D = UT + \frac{1}{2} aT^2$ *	A1 *
	(4)

Question 7

(i)(ii)	Using a correct strategy for solving the problem by setting up two equations in a and u only and solving for either	M1
	Equation in a and u only	M1
	$22 = 2u + \frac{1}{2} a 2^2$	A1
	Another equation in a and u only	M1
	$126 = 6u + \frac{1}{2} a 6^2$	A1
	5 m s^{-2}	A1
	6 m s^{-1}	A1ft

(7 m)