

Year 1 – Week 14 Exam Questions

Mark Scheme

Question 1

| Question | | Answer | Mks |
|----------|------|--|-----------------------|
| 1 | (i) | DR $(\sqrt{3})^7$ or $\sqrt{3^7}$ or $3^3 \times \sqrt{3}$ or $3\sqrt{243}$ $27\sqrt{3}$ | M1 A1 [2] |
| 1 | (ii) | DR $\frac{\sqrt{2}}{1-\sqrt{2}} \times \frac{1+\sqrt{2}}{1+\sqrt{2}}$ $= \frac{\sqrt{2}+2}{1-2}$ or $\frac{\sqrt{2}+2}{-1}$ or $\frac{\sqrt{2}+2}{1+\sqrt{2}-\sqrt{2}-2}$ $= -2 - \sqrt{2}$ ISW | M1 A1 A1 [3] |

Question 2

| | | | |
|---|------|--|---------------------|
| 2 | (i) | $3^2 - 4k = 0$ $k = \frac{9}{4}$ or 2.25 | M1 A1 [2] |
| 2 | (ii) | $(3-x)(2+x) > 0$ or $(x-3)(x+2) < 0$ $-2 < x < 3$ or $3 > x > -2$ ISW or $x \in (-2, 3)$ | M1 A1 [2] |

Question 3

| Question | | Answer | Mks |
|----------|------|--|-----------------------|
| 10 | (i) | $\frac{3}{8} + \frac{5}{16} + 4p + p = 1$ $p = \frac{1}{16}$ or 0.0625 | M1 A1 [2] |
| 10 | (ii) | $\frac{3}{8} \times \frac{5}{8}$ or $\frac{3}{8} \times \frac{3}{8}$ seen oe $\frac{3}{8} \times \frac{5}{8} + \frac{5}{8} \times \frac{3}{8} + \frac{3}{8} \times \frac{3}{8}$ oe $= \frac{39}{64}$ or 0.609 (3 sf) | M1 M1 A1 [3] |

Question 4

| Question | | Answer | Marks |
|----------|------|--|--|
| 1 | (i) | $\frac{\sin x}{20} = \frac{\sin 45}{16}$ $\sin x = \frac{20 \sin 45}{16} \left(= \frac{5\sqrt{2}}{8} \right)$ 62.1 and 117.9 | M1* A1 Dep*M1 A1 [4] |
| 1 | (ii) | $\frac{1}{2}(BC)(20)\sin(45) = 75\sqrt{2}$ (BC \Rightarrow) 15 (cm) | M1 A1 [2] |

Question 5

| Question | | Answer | Marks |
|----------|------|---|-----------------------------|
| 2 | (i) | $\frac{2}{3+x-4} \text{ or } \frac{2}{3+x+4}$ $y = \frac{2}{x-1}$ | M1 A1 [2] |
| | (ii) | Stretch Scale factor $\frac{5}{2}$ parallel to the y -axis | B1 B1 [2] |

Question 6

| Question | | Answer | Marks |
|----------|-------|---|--------------------------|
| 4 | (i) | $4[x^2 - 3x] + 11$ | |
| | | $4\left[\left(x - \frac{3}{2}\right)^2 - \frac{9}{4}\right] + 11$ | $a = 4$ |
| | | $4\left(x - \frac{3}{2}\right)^2 + 2$ | $(x - 3/2)^2$ $c = 2$ |
| | (ii) | No real roots | B1 [1] |
| | (iii) | $r = 0 \Rightarrow 1$ real root or 1 repeated root $r < 0 \Rightarrow 2$ real roots $r > 0 \Rightarrow$ no real roots | M1 A1 [2] |

Question 7

| Question | | Answer | Mks |
|----------|---------|---|--|
| 7 | (i) (b) | $a + \frac{1}{2}(c - a)$ or $c + \frac{1}{2}(a - c)$ | M1 |
| | | $= \frac{1}{2}(a + c)$ or $\frac{1}{2}a + \frac{1}{2}c$ | A1 [2] |
| | (ii) | $\vec{OB} = (a + c)$ $\Rightarrow \vec{OP} = \frac{1}{2}\vec{OB}$ Must see previous line $\Rightarrow P$ is midpt of OB or OPB is a straight line and $OP = PB$ Hence diagonals of //m bisect one another | M1 A1+ dep+ A1 E1 [4] |

Question 8

| Question | | Answer | Marks | AOs | Guidance |
|----------|---------|--|-----------------------------|----------------------------|--|
| 11 | (i) (a) | $18 = \left(\frac{8+u}{2}\right)(9)$ $u = -4$ therefore the speed of P is $4 \text{ (ms}^{-1}\text{)}$ | M1 A1 [2] | 3.4 1.1 | Use of $s = \left(\frac{u+v}{2}\right)t$ AG |
| | (i) (b) | eg $8 = -4 + 9a$ $a = \frac{4}{3} \text{ (ms}^{-2}\text{)}$ | M1 A1 [2] | 3.4 1.1 | Use of $v = u + at$ with their u or $s = vt - \frac{1}{2}at^2$ or $v^2 = u^2 + 2as$ with their u or $s = ut + \frac{1}{2}at^2$ with their u Accept 1.33 or better |
| Question | | Answer | Marks | AOs | Guidance |
| | (ii) | $0 = -4 + \frac{4}{3}t$ $t = 3$ $-s_{\max} = -4t + \frac{1}{2}\left(\frac{4}{3}\right)t^2$ $s_{\max} = 6 < 10$ so P is never at B | M1 A1 M1 A1 [4] | 3.1b 1.1 3.4 2.2a | Use of $v = u + at$ with $v = 0$ and their a and u Use of $s = ut + \frac{1}{2}at^2$ with their a, u & t Compare with 10 or suitable comment |
| | OR | $-10 = -4t + \frac{1}{2}\left(\frac{4}{3}\right)t^2$ e.g. det = -24 therefore not possible | M1 A1 | | Use of $s = ut + \frac{1}{2}at^2$ with their u and a and suitable s Consider $b^2 - 4ac$ or attempt to solve three term quadratic in t Or $36 - 60 < 0$ therefore not possible |
| | OR | $0 = (\pm 4)^2 + 2\left(\frac{4}{3}\right)s$ or $v^2 = (\pm 4)^2 + 2\left(\frac{4}{3}\right)(-10)$ $s = -6$ or $v^2 = -\frac{32}{3}$ Suitable conclusion | M2 A1 A1 | | Use of $v^2 = u^2 + 2as$ with their a and u and either $v = 0$ or $s = \pm 10$ Dependent on previous A mark |